

QTL reading list -- EDG Spring 2011

Ian and Jeff's suggestions:

- Abiola, O., J. M. Angel, P. Avner, A. A. Bachmanov, J. K. Belknap, B. Bennett, E. P. Blankenhorn et al. 2003. The nature and identification of quantitative trait loci: a community's view. Nature Reviews Genetics 4:911-916.**
- Barton, N. H., and P. D. Keightley. 2002. Understanding quantitative genetic variation. Nature Reviews Genetics 3:11-21.**
- Beraldi, D., A. F. McRae, J. Gratten, J. Slate, P. M. Visscher, and J. M. Pemberton. 2007. Mapping quantitative trait loci underlying fitness-related traits in a free-living sheep population. *Evolution* 61:1403-1416.
- Bergelson, J., and F. Roux. 2010. Towards identifying genes underlying ecologically relevant traits in *Arabidopsis thaliana*. Nature Reviews Genetics 11:867-879.**
- Blackman, B. K., J. L. Strasburg, A. R. Raduski, S. D. Michaels, and L. H. Rieseberg. 2010. The Role of Recently Derived FT Paralogs in Sunflower Domestication. *Current Biology* 20:629-635.
- Bouck, A., S. R. Wessler, and M. L. Arnold. 2007. Qtl analysis of floral traits in Louisiana Iris hybrids. *Evolution* 61:2308-2319.
- Buckler, E. S., J. B. Holland, P. J. Bradbury, C. B. Acharya, P. J. Brown, C. Browne, E. Ersoz et al. 2009. The Genetic Architecture of Maize Flowering Time. *Science* 325:714-718.
- Chen, K. Y., B. Cong, R. Wing, J. Vrebalov, and S. D. Tanksley. 2007. Changes in regulation of a transcription factor lead to autogamy in cultivated tomatoes. *Science* 318:643-645.
- Doerge, R. W. 2002. Mapping and analysis of quantitative trait loci in experimental populations. *Nature Reviews Genetics* 3:43-52.
- Fishman, L., A. J. Kelly, and J. H. Willis. 2002. Minor quantitative trait loci underlie floral traits associated with mating system divergence in *Mimulus*. *Evolution* 56:2138-2155.
- Flint, J., and T. F. C. Mackay. 2009. Genetic architecture of quantitative traits in mice, flies, and humans. Genome Research 19:723-733.**
- Flint, J., W. Valdar, S. Shifman, and R. Mott. 2005. Strategies for mapping and cloning quantitative trait genes in rodents. *Nature Reviews Genetics* 6:271-286.
- Gardner, K. M., and R. G. Latta. 2007. Shared quantitative trait loci underlying the genetic correlation between continuous traits. *Molecular Ecology* 16:4195-4209.
- Hall, M. C., C. J. Basten, and J. H. Willis. 2006. Pleiotropic Quantitative Trait Loci Contribute to Population Divergence in Traits Associated With Life-History Variation in *Mimulus guttatus*. *Genetics* 172:1829-1844.
- Kelly, J. K. 2009. Connecting QTLs to the G-matrix of evolutionary quantitative genetics. *Evolution* 63:813-825.
- Latta, R. G. 2009. Testing for local adaptation in *Avena barbata*: a classic example of ecotypic divergence. Molecular Ecology 18:3781-3791.**

- Mackay, T. F. C. 2001a. Quantitative trait loci in *Drosophila*. *Nature Reviews Genetics* 2:11-20.
- . 2001b. THE GENETIC ARCHITECTURE OF QUANTITATIVE TRAITS. *Annual Review of Genetics* 35:303-339.
- Mackay, T. F. C., E. A. Stone, and J. F. Ayroles. 2009. The genetics of quantitative traits: challenges and prospects. *Nat Rev Genet* 10:565-577.**
- Mauricio, R. 2001. Mapping quantitative trait loci in plants: uses and caveats for evolutionary biology. *Nature Trends Genetics* 2:370-381.
- McCarthy, M. I., G. R. Abecasis, L. R. Cardon, D. B. Goldstein, J. Little, J. P. A. Ioannidis, and J. N. Hirschhorn. 2008. Genome-wide association studies for complex traits: consensus, uncertainty and challenges. *Nature Reviews Genetics* 9:356-369.
- Moyle, L. C., and T. Nakazato. 2010. Hybrid Incompatibility "Snowballs" Between *Solanum* Species. *Science* 329:1521-1523.
- Paterson, A. H., E. S. Lander, J. D. Hewitt, S. Peterson, S. E. Lincoln, and S. D. Tanksley. 1988. Resolution of quantitative traits into Mendelian factors by using a complete linkage map of restriction fragment length polymorphisms. *Nature* 335:721-726.**
- Smith, M. W., and S. J. O'Brien. 2005. Mapping by admixture linkage disequilibrium: advances, limitations and guidelines. *Nat Rev Genet* 6:623-632.
- Wu, R. L., and M. Lin. 2006. Opinion - Functional mapping - how to map and study the genetic architecture of dynamic complex traits. *Nature Reviews Genetics* 7:229-237.**

Doug's suggestions:

- Beckmann J.S. and Soller M. (1983), Restriction fragment length polymorphisms in genetic improvement: methodologies, mapping and costs. *Theoretical and Applied Genetics* 67 p35-43.
- Botstein D., White R.L., Skolnick M. and Davis R.W. (1980), Construction of a genetic linkage map in man using restriction fragment length polymorphisms. *American Journal of Human Genetics* 32 p314-331.
- Breese E.L. and Mather K. (1957), The organization of polygenic activity within a chromosome in *Drosophila*. I. Hair characters. *Heredity* 11 p373-395.
- Carbonell E.A., Gerig T.M., Balansard E. and Asins M.J. (1992), Interval mapping in the analysis of non-additive quantitative trait loci. *Biometrics* 48 p305-315.
- Cowen N.M. (1988), The use of replicated progenies in marker-based mapping of QTLs. *Theoretical and Applied Genetics* 75 p857-862.

- Doganlar S., Tanksley S.D. and Mutschler M.A. (2000), Identification and molecular mapping of loci controlling fruit ripening time in tomato. *Theoretical and Applied Genetics* 100 p249-255.
- Fulton T.M., Beck-Bunn T., Emmatty D., Eshed Y., Lopez J., Petiard V., Uhlig J., Zamir D. and Tanksley S.D. (1997), QTL analysis of an advanced backcross of *Lycopersicon peruvianum* to the cultivated tomato and comparisons with QTLs found in other wild species. *Theoretical and Applied Genetics* 95 p881-894.
- Flowers T.J., Koyama M.L., Flowers S.A., Sudhakar C., Singh K.P. and Yeo A.R. (2000), QTL: their place in engineering tolerance of rice to salinity. *Journal of Experimental Botany* 51 p99-106.
- Geldermann H. (1975), Investigations on inheritance of quantitative characters in animals by gene markers. I. Methods. *Theoretical and Applied Genetics* 46 p319-330.
- Haley C.S. and Knott S.A. (1992), A simple regression method for mapping quantitative trait loci in line crosses using flanking markers. *Heredity* 69 p315-324.
- Hayes P., Prehn D., Vivar H., Blake T., Comeau A., Henry I., Johnstone M., Jones B., Steffenson B., St. Pierre C.A. and Chen F. (1996), Multiple disease resistance loci and their relationship to agronomic and quality loci in a spring barley population. *Journal of Quantitative Trait Loci*.
- Hyne V. and Kearsey M.J. (1995), QTL analysis: further uses of 'marker regression'. *Theoretical and Applied Genetics* 91 p471-476.
- Jansen R.C. (1993), Interval mapping of multiple quantitative trait loci. *Genetics* 135 p205-211.
- Jansen R.C. (1996), A general monte carlo method for mapping multiple quantitative trait loci. *Genetics* 142 p305-311.
- Kao C-H., Zeng Z-B and Teasdale R.D. (1999), Multiple interval mapping for quantitative trait loci. *Genetics* 152 p1203-1216.
- Kearsey M.J. and Farquhar A.G.L. (1998), QTL analysis in plants; where are we now? *Heredity* 80 p134-142.
- Kearsey M.J. and Hyne V. (1994), QTL analysis a simple 'marker regression' approach. *Theoretical and Applied Genetics* 89 p698-702.
- Lander E.S. and Botstein D. (1989), Mapping mendelian factors underlying quantitative traits using RFLP linkage maps. *Genetics* 121 p185-199.
- Larkin J.C. Young N., Prigge M. and Marks M.D. (1996), The control of trichome spacing and number in *Arabidopsis*. *Development* 122 p997-1005.

- Long A.D., Mullaney S.L., Reid L.A., Fry J.D., Langley C.H. and Mackay T.F.C. (1995), High resolution mapping of genetic factors affecting abdominal bristle number in *Drosophila melanogaster*. *Genetics* 139 p1273-1291.
- Luo Z.W. and Kearsley M.J. (1992), Interval mapping of quantitative trait loci in an F2 population. *Heredity* 69 p236-242.
- Mangin B., Thoquet P., Olivier J. and Grimsley N.H. (1999), Temporal and multiple quantitative trait loci analyses of resistance to bacterial wilt in Tomato permit the resolution of like loci. *Genetics* 151 p1165-1172.
- Martinez O. and Curnow R.N. (1992), Estimating the locations and the sizes of effects of quantitative trait loci using regression mapping. *Theoretical and Applied Genetics* 85 p480-488.
- Nettleton D. and Praestgaard J. (1998), Interval mapping of quantitative trait loci through order-restricted inference. *Biometrics* 54 p74-87.
- Oberthur L., Blake T.K., Dyer W.E. and Ullrich S.E. (1995), Genetic analysis of seed dormancy in barley (*Hordeum vulgare L.*). *Journal of Quantitative Trait Loci*.
- Qi X. and Lindhout P. (1997), Development of AFLP markers in barley. *Molecular and General Genetics* 254 p330-336.
- Richter K., Schondelmaier J. and Jung C. (1998), Mapping of quantitative trait loci affecting *Drechslera teres* resistance in barley with molecular markers. *Theoretical and Applied Genetics* 97 p1225-1234.
- Rogot M. and Hoisington D.A. (1993), Molecular marker for plant breeding: comparisons of RFLP and RAPD genotyping costs. *Theoretical and Applied Genetics* 86 p975-984.
- Sax K. (1923), The association of size differences with seed-coat pattern and pigmentation in *Phaseolus vulgaris*. *Genetics* 8 p552-560.
- Swarup K., Alonso-Blanco C., Lynn J.R., Michaels S.D., Amasino R.M., Koorneef M. and Millar A.J. (1999), Natural allelic variation identifies new genes in the *Arabidopsis* circadian system. *The Plant Journal* 20 p67-77.
- Tanksley S. (1993), Mapping polygenes. *Annual review of Genetics* 27 p205-233.
- Thoday J.M. (1961), Location of polygenes. *Nature* 191 p368-370.
- Whittaker J.C., Thompson R. and Visscher P.M. (1996), On the mapping of QTL by regression of phenotypes on marker type. *Heredity* 77 p23-32.

- Wright A.J. and Mowers R.P. (1994), Multiple regression for molecular-marker, quantitative trait data from large F2 populations. *Theoretical and Applied Genetics* 89 p305-312.
- Wu W.R. and Li W.M. (1994), A new approach for mapping quantitative trait loci using complete genetic marker linkage maps. *Theoretical and Applied Genetics* 89 p535-539.
- Wu W.R. and Li W.M. (1996), Model fitting and model testing in the method of joint mapping of quantitative trait loci. *Theoretical and Applied Genetics* 92 p477-482.
- Xu S. (1995), A comment on the simple regression method for interval mapping. *Genetics* 141 p1657-1659.
- Xu S. (1998), Further investigation on the regression method of mapping quantitative trait loci. *Heredity* 80 p364-373.
- Yamamoto T., Lin H., Sasaki T. and Yano M. (2000), Identification of heading date quantitative trait locus *Hd6* and characterisation of its epistatic interactions with *Hd2* in rice using advanced backcross progeny. *Genetics* 154 p885-891.
- Yan J.Q., Zhu J., He C.X., Benmoussa M. and Wu P. (1998), Quantitative trait loci analysis for the developmental behaviour of tiller number in rice (*Oryza sativa L.*). *Theoretical and Applied Genetics* 97 p267-274.
- Yin X., Stam P., Johan Dourleijn C. and Kropff M.J. (1999), AFLP mapping of quantitative trait loci for yield-determining physiological characters in spring barley. *Theoretical and Applied Genetics* 99 p244-253.
- Zeng Z-B. (1994), Precision mapping of quantitative trait loci. *Genetics* 136 p1457-1468.

Barry's suggestions:

Association studies:

Lango Allen H, Estrada K, Lettre G, Berndt S. I, Weedon M. N, et al. (2010) Hundreds of variants clustered in genomic loci and biological pathways affect human height. *Nature* 467: 832–838.

Manolio T. A, Collins F. S, Cox N. J, Goldstein D. B, Hindorff L. A, et al. (2009) Finding the missing heritability of complex diseases. *Nature* 461: 747–753.

Goldstein DB. The importance of synthetic associations will only be resolved empirically. *PLoS Biol.* 2011 Jan 18;9(1):e1001008.

Wang K, Dickson S. P, Stolle C. A, Krantz I. D, Goldstein D. B, et al. (2010) Interpretation of association signals and identification of causal variants from genome-wide association studies. *Am J Hum Genet* 86: 730–742.

Goldstein DB, 2011 The Importance of Synthetic Associations Will Only Be Resolved Empirically. PLoS Biol 9(1): e1001008.

Dickson S. P, Wang K, Krantz I, Hakonarson H, Goldstein D. B (2010) Rare variants create synthetic genome-wide associations. PLoS Biol 8: e1000294.

Anderson CA, Soranzo N, Zeggini E, Barrett JC, 2011 Synthetic Associations Are Unlikely to Account for Many Common Disease Genome-Wide Association Signals. PLoS Biol 9(1): e1000580.

Wray NR, Purcell SM, Visscher PM, 2011 Synthetic Associations Created by Rare Variants Do Not Explain Most GWAS Results. PLoS Biol 9(1): e1000579.

Shields R, 2011 Common Disease: Are Causative Alleles Common or Rare?. PLoS Biol 9(1): e1001009.

Eichler EE, Flint J, Gibson G, Kong A, Leal SM, Moore JH, Nadeau JH. Missing heritability and strategies for finding the underlying causes of complex disease. Nat Rev Genet. 2010 Jun;11(6):446-50.

QTL mapping:

Kroymann J, Mitchell-Olds T. Epistasis and balanced polymorphism influencing complex trait variation. Nature. 2005 May 5;435(7038):95-8.

Gerke J, Lorenz K, Ramnarine S, Cohen B. Gene-environment interactions at nucleotide resolution. PLoS Genet. 2010 Sep 30;6(9).

Gerke J, Lorenz K, Cohen B. Genetic interactions between transcription factors cause natural variation in yeast. Science. 2009 Jan 23;323(5913):498-501.

Deutschbauer AM, Davis RW. Quantitative trait loci mapped to single-nucleotide resolution in yeast. Nat Genet. 2005 Dec;37(12):1333-40.

Ehrenreich IM, Torabi N, Jia Y, Kent J, Martis S, Shapiro JA, Gresham D, Caudy AA, Kruglyak L. Dissection of genetically complex traits with extremely large pools of yeast segregants. Nature. 2010 Apr 15;464(7291):1039-42.

Ehrenreich IM, Gerke JP, Kruglyak L. Genetic dissection of complex traits in yeast: insights from studies of gene expression and other phenotypes in the BYxRM cross. Cold Spring Harb Symp Quant Biol. 2009;74:145-53.

Cubillos FA, Billi E, Zörgö E, Parts L, Fargier P, Omholt S, Blomberg A, Warringer J, Louis EJ, Liti G. Assessing the complex architecture of polygenic traits in diverged yeast populations. Mol Ecol. 2011 Jan 25.

Kim HS, Fay JC. A combined-cross analysis reveals genes with drug-specific and background-dependent effects on drug sensitivity in *Saccharomyces cerevisiae*. *Genetics*. 2009 Nov;183(3):1141-51.